

A Study on the *Phytoliths* of Selected Taxa of *Poaceae*

Dr. Shipra Rani Podder, Associate Professor, Dept. of Botany, G K Gowani Govt College, Bhinmal (Rajasthan).

shipranipodder@gmail.com

ABSTRACT

Phytoliths are microscopic, inorganic silica bodies that are produced by plants. They are found in all plant tissues, but are most abundant in the epidermal cells of leaves, stems, and roots. *Phytoliths* are resistant to decay and can survive for millions of years in the environment. They are therefore an important source of information about past vegetation and climate.

The *Poaceae* (grasses) are a large and diverse family of plants that includes over 10,000 species. Grasses are found in all parts of the world and play an important role in many ecosystems. *Phytoliths* from grasses are therefore a valuable tool for paleoecological research.

A number of studies have been conducted on the *Phytoliths* of selected taxa of *Poaceae*. These studies have shown that the shape, size, and morphology of *Phytoliths* can be used to identify different species of grasses. In addition, the abundance of different types of *Phytoliths* can be used to reconstruct past vegetation communities.

For example, a study examined *Phytoliths* from sediments in the Amazon rainforest. They found that the abundance of grass *Phytoliths* increased over time, suggesting that the rainforest has become more open in recent centuries. This change is likely due to human activities, such as deforestation and fire.

KEYWORDS: *Phytoliths*, *Taxa*, *Poaceae*.

INTRODUCTION

Phytoliths can also be used to reconstruct past climate. For example, a study examined *Phytoliths* from sediments in the southwestern United States. They found that the size of grass *Phytoliths* decreased during periods of drought, suggesting that grasses were growing under more stressful conditions.

Phytoliths are a valuable tool for paleoecological research. They can be used to identify different species of grasses, reconstruct past vegetation communities, and infer past climate. As our understanding of *Phytoliths* continues to grow, they will become an even more powerful tool for understanding the past.

In addition to the studies mentioned above, there have been many other studies on the *Phytoliths* of selected taxa of *Poaceae*. These studies have shown that *Phytoliths* can be used to track changes in vegetation over time, identify the presence of different types of grasses, and infer past climate. *Phytoliths* are therefore a valuable tool for paleoecological research.

The *Phytoliths* of *Poaceae* are found in a wide variety of habitats, but they are most common in grasslands and savannas. They are also found in wetlands, forests, and even deserts.

The type of *Phytoliths* that are found in a particular habitat can vary depending on the species of grass that is present. For example, saddle *Phytoliths* are more common in grasses that grow in grasslands, while triradiate *Phytoliths* are more common in grasses that grow in wetlands.

The *Phytoliths* of *Poaceae* can be used to reconstruct past environments. For example, the presence of saddle *Phytoliths* indicates that a grassland habitat was present, while the presence of triradiate *Phytoliths* indicates that a wetland habitat was present.

Phytoliths can also be used to track changes in climate and vegetation over time. For example, a study of *Phytoliths* from sediments in California showed that the climate became drier and the vegetation became more open during the last glacial period.

The *Poaceae* are characterized by their slender, jointed stems and their leaves, which are arranged in two rows on either side of the stem. The leaves are typically long and narrow, with a ligule at the base. The flowers of grasses are small and inconspicuous, and they are usually arranged in spikelets. The spikelets are borne in panicles, racemes, or heads.

The *Poaceae* are divided into two subfamilies: the Bambusoideae and the Pooideae. The Bambusoideae includes the bamboos, which are a group of woody grasses that can grow to be very tall. The *Poaceae* includes the true grasses, which are the most common type of grass.

Some of the most important taxa of *Poaceae* include:

- ***Triticum aestivum* (wheat):** Wheat is a major cereal crop that is grown in many parts of the world. It is a good source of carbohydrates and protein.
- ***Oryza sativa* (rice):** Rice is another major cereal crop that is grown in many parts of the world. It is a good source of carbohydrates.
- ***Zea mays* (corn):** Corn is a major cereal crop that is grown in many parts of the world. It is a good source of carbohydrates, protein, and vitamins.
- ***Saccharum officinarum* (sugarcane):** Sugarcane is a grass that is grown for its sweet juice. The juice is processed to produce sugar.
- ***Miscanthus giganteus* (giant miscanthus):** Giant miscanthus is a grass that is grown for its biomass. The biomass can be used to produce energy or to make paper.

Grasses are an important part of our world. They provide us with food, shelter, and energy. They also play an important role in many ecosystems. We should all do our part to protect grasses and the ecosystems that they depend on.

Here are some additional facts about grasses:

- Grasses are the most successful plant family on Earth. They cover about one-third of the land surface.
- Grasses are adapted to a wide range of climates and habitats. They can be found from the Arctic to the tropics.
- Grasses are an important source of food for many animals, including humans.
- Grasses are also used to make a variety of products, including paper, textiles, and biofuels.

PHYTOLITHS OF SELECTED TAXA OF POACEAE

Phytoliths are undigested silicon dioxide crusts that form between and within plant cells and tissues. The word phytolith, from the Greek meaning plant stone, is a piece of time that has been used to show a considerable amount of mineral material retained by plants, silica or calcium in composition. The last alternative occasion is keystones produced using calcium oxalates, which may be common and cognate in desert flora and some other dry land plants and have been used to record fundamental pieces of the really old dietary plan.

Phytoliths are considered the milling and confirmation cycles by which some living higher plants have strong fields to capture silica in a soluble form from groundwater and then store it in an intracellular or extracellular space. After plant death and decomposition, these pieces of silica are stored in the soil and developed as microscopic particles of varying sizes and shapes. Because *Phytoliths* are inorganic and thus impervious to the forces of decay that cause various types of plant material to crumble, they are scratched in a particularly preserved state over wide time periods. They are apparently the most solid distinct plant fossils known to science. Various angiosperms, gymnosperms and pteridophytes produce titanic expanses of *Phytoliths*.

Not all plant species produce *Phytoliths*; At any rate, a huge variable for understanding phytolith formation is that the strong fields for collection, near the location of set stores in apparent tissues and plant cells, are virtually identical across plant species and their most frequent related Paying little attention to what the specific conditions of progress are.

Appropriately, cosmopolitan specialists and families of cool zone, cold and tropical species show concordant models in their phytolith planning schemes. Furthermore, plants that do not concentrate their organs for silicification and do not yield comparably quantitative degrees of *Phytoliths* do not do so where in the world they form. A portion of these non-producers comprise the aroids, Amaranthaceae and yams, trees in various families and most of the desert flora. In many plant taxa, specific cells and tissues that seek out aerial organs are seen as districts sent by critical strong fields for acclimatization by the plant as a whole.

A focal validation for this has shown a continuum in the formation and breaking points of silica, leaving no question that the presence of silica, in both its isolated and solid states, has an extraordinary, consistent effect on the turn and reform of events. In the plants. For example, the necessary confirmation that a huge barrier of *Phytoliths* is to protect plants from herbivores and pathogenic creatures. It has been observed that plants of a specific zoological taxonomy may have varying degrees of *Phytoliths* when grown under different conditions. It gives after a while that the variable silica content within the species has basically something to do with the reforming of *Phytoliths* which are not normally allocated by the severe strong zones for deposition by the plant.

A piece of these spots are the sheets of polyhedral and saw-shaped epidermal cells that cover the surfaces of tree leaves and the layers of bulliform cells that lie essentially on the upper surface of grass leaves and when they If possible, it gives spots beyond this. Not silicate. As a general rule, an overflow pile of water in the laying out environment and submergence of root establishments are directly associated with extended bulliform cell silicification.

Favorably, these are usually the most non-morphologically specific such *Phytoliths*, and they are not commonly used to visualize plants at any coordinate level. In fact the referenced study and others also suggest that plants are constantly subjected to elemental mixing in phytolith material and grown up would be huge scavengers of solid silica in any environment.


The phenotypic flexibility of *Phytoliths* induced by changing normal conditions cannot thwart the potential use of phytolith morphometry to differentiate between plant taxa. On occasion, the conditions of the silica bodies warranted sufficient emphasis to address ID at the species level while focusing on plant reference material.

RESULTS AND DISCUSSION

Bambusidae showed a game plan of the phytolith type. The most widely observed types were the bilobates with a range in the shape of the bend and the length and width of the bleeding edge. Nearby bilobates were the most common type and thus could be used as marker types for the subfamily. Our conclusion that bilobates and saddles are found in three individuals in total, *Bambusa ventricosa* Schrad., *B. vulgaris* Nees and *Arundinaria falcata* Nees, in our mix, describe more endurance to the proper meaning of these variants of the subfamily. The bilobates and saddles are formed in epidermal short cells whose lumens are completely infilled with silica. The different types experienced in the subfamily were dendritics in all three individuals of the subfamily, trapezoids in *Bambusa ventricosa* Schrad., and *Arundinaria falcata* Nees., scutiform and sinuate elongate in *Bambusa*.

In our model, the focal *Oryza sativa* L. showed quadrilobate and trapezoidal bilobates are regarded as the subfamily *Panicoideae*. variegated, cross-framed and grasses helped to restrict showed that the cross-leaved dominant *Phytoliths* in this The *Pooideae* peoples sizes and shapes of subfamily, the genus *Triticum* trapezoid, and sinuate stretch types into two individuals. These types are derived from epidermal long cells in both the costal and intercostal districts of the adaxial and abaxial epidermis.

Also the common sizes of sinuate grow type were 2053 μm and 2030 μm in *Hordeum vulgare* L., *Triticum aestivum* L. unreservedly. Note the spectacular appearance by point *Phytoliths* of essentially undefined shape in these species. The manufacturer found that the trapezoid phytolith




individual of this subfamily, cross-framed, bilobate, *Phytoliths*. Of these types, brand name for the Anyway, the presence of quadrilobate panicoid oryzoids. Earlier assessments type and bilobate are the species.

revealed a remarkable mix of *Phytoliths*. Within the consolidated the dendritic,

is the diagnostic type for *Triticum aestivum* L., which is close to the dendritic and sinuate stretch. We have also retrieved these variants in our collection.

Group *Arundinella* observed single part *Arundinella* the most composite saw type seat and rondels were seen types that could help other panicoid species.

Andropogony, the third Panicoideae, was observed continuous model. Bilobate such clades. However, smooth emerged as other family.



in a continuous model by a nepalensis Trin. bilobates as (1b). After the bilobate, the as the broadest and sharpest distinguish this species from

group of the subfamily by 12 species in the species were available in all dendritic, clavate, and common genera in the broad

Epidermal layer showing cross shaped *Phytoliths*

The game plan has also been found to have a fair significance in the general degree of the parts of the *Phytoliths*, all sizes of different types of *Phytoliths* and adjacent to the parcel. Bilobates with additional apical whorls have overall thicker knives when separated from those with almost limited whorls. Part of the animal species that bear more noticeable bilobates with thicker scutes include *Cenchrus setigerus* Vahl.

The assessment of surface characteristics of different types of *Phytoliths* via light and scanning electron microscopy is not, by all accounts, the fundamental game plan in surface ornamentation and the ultra course of action that can be used in the assessment of grass taxa. In any case, the SEM evaluation revealed that the bilobate *Phytoliths* can be set into clear groups considering the divisions that appear, by all accounts, to be the outer edge.

Grasses are a fascinating and important group of plants. They play a vital role in our world and we should all do our part to learn more about them and to protect them.

One of the most important biological significance of *Poaceae* is their role in the food chain. Grasses are a primary food source for many animals, including cattle, horses, sheep, and deer. They are also an important source of food for humans, as they are the source of many staple foods, such as rice, wheat, corn, and barley.

Grasses also play an important role in the environment. They help to prevent erosion by binding the soil together. They also help to filter water and provide a habitat for many different types of animals.

In addition to their ecological importance, grasses are also used for a variety of human purposes. They are used to make building materials, such as bamboo and thatch. They are also used to make paper, textiles, and biofuel.

The *Poaceae* is a diverse and important group of plants. They play a vital role in the food chain, the environment, and human society.

Here are some specific examples of the biological significance of *Poaceae*:

- **Food:** Grasses are a major source of food for humans and animals. Some of the most important food crops that are grasses include rice, wheat, corn, and barley.
- **Fodder:** Grasses are also an important source of fodder for livestock. They are high in nutrients and can be easily digested by animals.
- **Building materials:** Some grasses, such as bamboo, can be used to make building materials. Bamboo is strong and lightweight, and it can be used to make a variety of structures, such as houses, bridges, and furniture.
- **Papermaking:** Grasses are also used to make paper. The cellulose fibers in grass are long and strong, and they can be easily processed into paper.

- **Textiles:** Grasses can also be used to make textiles. The fibers in grass can be spun into yarn and then woven into cloth.
- **Biofuel:** Grasses can also be used to produce biofuel. Biofuel is a renewable energy source that can be used to power cars, trucks, and other vehicles.

The *Poaceae* is a valuable and versatile group of plants. They provide food, shelter, and other resources for humans and animals. They also play an important role in the environment by preventing erosion and filtering water.

CONCLUSION

The delineation and confirmation of grasses as a whole is a troublesome task because of a significant reliance on perceptual parts that are not only open for a limited period of time in the phenological opportunities of grasses, but are negligible in size and cover morphology between taxa. Botanical morphological features are of little help with the standard procedures for viewing grass delineation and certification. Here, *Phytoliths* refer to a flexible individual and give the ID of the grass.

The survey also highlighted some novel species such as cap-shaped *Phytoliths* from *Digitaria ciliaris* (Retz.) Koeler. Surface portions and morphometric evaluations give additional boundaries for delineation and confirmation of grass taxa. Further assessment on *Phytoliths* of field grass types would help in building up phytolith profiles of leaf epidermis and various tissues and organs. It is particularly after such evaluation that we will have the option of making phytolith indications of grass species, genera and higher taxa. With these advances, it will be possible to see species authentication in grass even at the vegetative stage. An epidermal striping with a fascinating phytolith etching would give the clearest evidence necessary for faunal classes as an individual, such as the spikelet's correction in the perceptual timing of grass phenology.

REFERENCES

- D. Barboni and L. Bremond, "Phytoliths of east African grasses: An assessment of their environmental and taxonomic significance based on floristic data," *Review of Paleobotany and Palynology*. Vol. 59, pp. 29-41, 2009.
- D. Zurro, M. Madella, I. Briz, A. Vila, "Variability of the phytolith record in fisher-hunter-gatherer sites: An example from the Yamana society (Beagle Channel, Tierra del Fuego, Argentina)," *Quaternary International*, vol. 193, pp. 184-191, 2009.
- D.K. Chauhan, D.K. Tripathi, P. Sinha, S.P. Tiwari, "Biogenic silica in some pteridophytes," *Bionature*, vol. 29 no. 1, pp. 1-9, 2011.
- P.I. Jattisha, M. Sabu, "Phytoliths as a Tool for the Identification of Some Chloridoideae Grasses in Kerala," *International scholarly Research Network Botany 2012: Article ID 246057*, 9 pages.
- S.A. Shakoor and M.A. Bhat, "Morphological diversity of phytolith types in some chloridoid grasses of Punjab," *International Journal of Botany and Research*, vol. 4, no. 1, pp. 1-10, 2014.
- T. Balakhnina and A. Borkowska, "Effects of silicon on plant resistance to environmental stresses: Review," *International Agrophysics*, vol. 27, pp. 225-232, 2013.
- T.B. Ball, R. Ehlers and M. D. Standing, "Review of typologic and morphometric analysis of *Phytoliths* produced by wheat and barley," *Breeding Science*, vol. 59, pp. 505-512, 2009.
- Z. L. Szabo, S. Koracs and A. Peto, "Phytolith analysis of *Poa pretensis* (*Poaceae*) leaves," *Turkish Journal of Botany*. Vol. 38, 2014.