



Mathematical Modelling on Nutrients Quality in the Ground Water through Comparison of Abohar and Sri Ganganagar Places

Dr. Swatmaram, Associate Professor, Chaitanya Bharathi Institute of Technology, Hyderabad.

ABSTRACT

It is glowing notorious that agriculture is main occupation of citizens of Sri Ganganagar and Abohar cities and their closely villages. As insufficient water supply through rivers and canals is conjoint snag in these areas of Punjab and Rajasthan, So that farmers are customarily dependent on ground-water and cope with lack of water, they are using to dig tube glowings in farms for the growth of crops should not be affected. The main factor behind this is nutrient present in ground water, because these get dissolved in the soil, then through crops nutrients and minerals enter into the human body by Vegetables and Grains. It is fact that healthy crops give rise to healthy human-being.

In this research, we will collect such data from various farms and after Collection of data from different location, will be analyzed in order to compare the amount of nutrients and chemical present in the ground water. Here we are presenting the merits & demerits of irrigation using ground water through the Mathematical Modelling and comparative study.

Main objective of study to compare nutrients (which makes human healthy or unhealthy) and chemicals (which are harmful for health), their level of presence and their impact on human-being. The whole study is focused to prepare a model, to know the variation of nutrients of ground water from one place to another place. We recognize the scenario of ground water flow and level of nutrients after comparison.

Keywords: Modeling, Water, Soil, Nutrients, Mathematical.

Introduction

Mathematical Modelling:

Mathematical models can take sundry schemes, including vibrantal schemes, geometric models, disparity calculations, or spirited conjectural models. These and other types of models can overlap, with a given model involving a variety of abstract structures. In general, mathematical models may include logical models. In many cases, the quality of a precise field depends on how glowing the mathematical models developed on the conjectural side settle with fallouts of repeatable experiments. Lack of settlement between conjectural mathematical models and investigational measurements often leads to important advances as better theories are developed.

In the physical sciences, a traditional mathematical model contains most of the following elements:

1. Overriding reckonings
2. Auxiliary sub-models
3. Outlining reckonings
4. Constitutive reckonings
5. Expectations and restrictions
6. Preliminary and periphery situations
7. Classical restrictions and kinematic reckonings

Classifications:

Mathematical models are customarily unruffled of connections and variables. Connections can be described by machinists, such as algebraic machinists, utilities, differential machinists, etc. Variables are abstractions of scheme parameters of interest that must be quantified. Several classification criteria can be used for mathematical models according to their structure:

- **Linear vs. Non-Linear:** If all the machinists in a mathematical model unveil linearity, the flouting mathematical model is demarcated as linear. A model is considered to be non-linear



otherwise. The delineation of linearity and non-linearity is dependent on context, and linear models may have non-linear jargons in them. For example, in a statistical linear model, it is assumed that a connection is linear in the parameters, but it may be non-linear in the predictor variables. Similarly, a differential reckoning is supposed to be linear if it can be written with linear differential machinists, but it can still have non-linear jargons in it. In a mathematical programming model, if the objective utilities and restrictions are represented entirely by linear reckonings, then the model is regarded as a linear model. If one or more of the objective utilities or restrictions are represented with a non-linear reckoning, then the model is notorious as a non-linear model.

Non-linearity, even in impartially simple schemes, is often associated with portents such as chaos and irreversibility. Although there are exceptions, non-linear schemes and models tend to be more demanding to study than linear ones. A conjoint tactic to non-linear snags is linearization, but this can be snag tic if one is trying to study aspects such as irreversibility, which are strongly tied to non-linearity.

- **Static vs. vibrant:** A vibrant model accounts for time-dependent changes in the state of the scheme, while a static model calculates the scheme in equipoise, and thus is time-invariant. Vibrant models archetypally are represented by differential reckonings or metamorphosis reckonings.
- **Explicit vs. implicit:** If all of the input parameters of the overall model are notorious, and the output parameters can be calculated by a finite series of computations, the model is supposed to be explicit. But sometimes it is the output parameters which are notorious, and the corresponding inputs must be solved for by an iterative procedure, such as Newton's method (if the model is linear) or Broyden's method (if non-linear). In such a case the model is supposed to be implicit. For example, a jet engine's physical properties such as turbine and nozzle throat areas can be explicitly calculated given a design thermovibrant cycle (air and fuel flow rates, pressures, and temperatures) at a specific flight situation and power setting, but the engine's operating cycles at other flight situations and power settings cannot be explicitly calculated from the constant physical properties.
- **Discrete vs. continuous:** A discrete model treats objects as discrete, such as the particles in a molecular model or the states in a statistical model; while a continuous model represents the objects in a continuous manner, such as the velocity field of fluid in pipe flows, temperatures and stresses in a solid, and electric field that applies continuously over the entire model due to a point charge.
- **Deterministic vs. probabilistic (stochastic):** A deterministic model is one in which every set of variable states is uniquely determined by parameters in the model and by sets of previous states of these variables; therefore, a deterministic model always performs the same way for a given set of Preliminary situations. Conversely, in a stochastic model—customarily called a "statistical model"—randomness is present, and variable states are not described by unique values, but rather by probability distributions.
- **Deductive, inductive, or floating:** A deductive model is a logical structure based on a theory. An inductive model arises from empirical findings and generalization from them. The floating model rests on neither theory nor observation, but is merely the invocation of expected structure. Application of mathematics in social sciences outside of economics has been criticized for unfounded models. Application of catastrophe theory in science has been characterized as a floating model.



Construction:

In business and engineering, mathematical models may be used to maximize a certain output. The scheme under consideration will require certain inputs. The scheme relating inputs to outputs depends on other variables too: decision variables, state variables, exogenous variables, and random variables.

Decision variables are sometimes notorious as independent variables. Exogenous variables are sometimes notorious as parameters or constants. The variables are not independent of each other as the state variables are dependent on the decision, input, random, and exogenous variables. Furthermore, the output variables are dependent on the state of the scheme (represented by the state variables).

Objectives and restrictions of the scheme and its users can be represented as utilities of the output variables or state variables. The objective utilities will depend on the perspective of the model's user. Depending on the context, an objective function is also notorious as an index of performance, as it is some measure of interest to the user. Although there is no limit to the number of objective utilities and restrictions a model can have, using or optimizing the model becomes more involved (computationally) as the number increases.

For example, economists often apply linear algebra when using input-output models. Complicated mathematical models that have many variables may be consolidated by use of vectors where one symbol represents several variables.

Review of Literature

Salt affected soils in Punjab occupy large areas in the south-west and central regions and vary widely in salt composition, internal drainage and pedogenic processes. Continued irrigation with sodic (RSC) ground water has enhanced salt enrichment leading to soil structural deterioration and poor crop productivity. Ten soil profiles were collected from old and recent alluvial plains covering seven districts and analyzed to assess precise reclamation and management options. In irrigated areas, waterlogging occurred in sandy alluvial plain (Pedin 1 and 2) showing loamy sand to sandy loam soil texture, low CEC and the significant contents of CaCO_3 calcretes (15.5%) at 0.5 m depth. The Na^+ , Cl^- , SO_4^{2-} and at places Ca^{2+} and Mg^{2+} salts were dominant and indicated the saline nature of soil. Sodic soils (Pedin 3 and 4) were located in the recent alluvial plain that showed high pHs (9.3 to 9.9), ESP (72.1 to 81.0) and higher contents of Na^+ , CO_3^{2-} and HCO_3^- salts. Sodic soils (Pedin 5 and 6) were also located in the old alluvial plain under canal irrigation, and showed high water table depth (2.0 m), high pHs (8.8 to 9.1), high surface salt content (EC_e 20.8 dS m^{-1}) and ESP (54.1 to 81.2). The blocky soil structure indicated impermeable soil strata and Na^+ saturation of soil matrix. At places, sodic soils were reclaimed (Pedin 7, 8 and 9) as evident from low surface pHs (8.6) and were being used for ricewheat crops. In the sub-surface layers, these soils showed higher pHs (9.0 to 10.2) and higher contents of Na^+ , CO_3^{2-} and HCO_3^- . Barren sodic soil (Pedin 10) was located in the old alluvial plain of central Punjab showing brackish ground water with high RSC. Sodic soils were also located in the Ghaggar plain of Patiala district showing moderate pHs (8.9 to 9.2), ESP (64 to 76), NaHCO_3 content (5.0 to 5.5 meqL $^{-1}$) and SAR (46 to 52). Suitable management options and alternate land uses were suggested for growing salt tolerant crops, horticulture and forestry plantations with proper water management practices.

About 7% of world's total arable land is classified as salt affected lands, while in India it is around 6.73 million hectare (mha). With the requirement for more food driving agriculture to



find new areas and methods for greater productivity, the reclamation of the barren salt affected lands attains utmost importance. Though there are multiple methods to manage and bring back such soils under cultivation, vegetative remediation is of interest, being more cost-effective and environment friendly. The other methods (leaching, organic/chemical amendments) have restrictions depending on type of soil, availability of water, chances for loss of soil nutrients and beneficial micro-organisms; in addition to being more expensive to farmers. Phytoremediation can be demarcated as the cultivation of salt-tolerant or accumulating (halophyte) plants for the reduction of soil salinity and/or sodicity. There are several halophytes which can be grown at very high levels of electrical conductivity. For examples *Eucalyptus occidentalis* and *E. sargentii* (useful landscape trees) can withstand salinity over 30 dS m⁻¹. About 290 tree species of economic importance have been documented which could tolerate salinity levels of 7 to 8 dS m⁻¹. These plants provide multiple benefits like

Impact of Agra canal and tube glowing water irrigation on crops grown in Agra district, Uttar Pradesh Soil salinity is a major snag of Gujarat state and 2.22-million-hectare area is salt affected. Ground water quality in the region is also poor having variable salinity and therefore, there is a scarcity of good quality water for agriculture. Millets are small-seeded grasses that are hardy and grow glowing in dry zones as rain-fed crops, under marginal soil fertility and limited moisture. They are also unique due to their short growing season so possibility can be explored to adopt these crops for cultivation on saline soils. Seed germination and early seedling growth are crucial periods for crop cycles under salt stress, and determine the survival of plants. A petriplate experiment was conducted during August 2015 at ICAR-Central Soil Salinity Research Institute, Regional Research Station, Bharuch to study germination of finger millet and little millet varieties under application of different levels of saline water. Experiment was laid out in petriplates with two crops; Finger millet (*Eleusine coracana* Gaertn.); three varieties - GN-4, GN-5 and GNN-6 and Little millet (*Panicum sumatrense*); two varieties - GV-1 and GV-2; with five salinity levels (Distill water; EC 4, 8 12 and 16 dSm⁻¹). Fifty seeds of each variety were germinated on filter paper in closed Petri dishes with different level of saline water using a completely randomized design with three replications. Germinated seeds were counted daily from each replication and the growth parameters were calculated on the 15th day. According to the method of Wang and Wang (2006), salt damage index (Salt damage index SDI % = (germination percentage under the control situation - germination percentage under salt stress)/ germination percentage under the control situation X 100) was adopted to evaluate the salt tolerance of these varieties. Fallouts revealed that at salinity level of 16 dSm⁻¹ no seed of any variety seed was germinated. At salinity level of 12 dSm⁻¹ germination of seed of all three variety of finger millet was severely affected and seed of both varieties of little millet did not germinate. Highest germination per cent (87 and 93 %) and lowest salt damage index (0.0 and 1.8 %) observed at salinity level of 4 dSm⁻¹ in finger millet (GNN-6) and little millet (GV-2) respectively. At salinity level of 8 dS m⁻¹ finger millet (GNN-6) and little millet (GV-2) showed maximum germination percentage (72 and 78 %) and lowest salt damage index (17.3 and 17.5%). Gradual reduction was observed at increased level of salinity upto 8 dS m⁻¹, and at 12 dS m⁻¹, severe reductions were observed for shoot length, root length and fresh & dry weight of seedlings in all varieties. These fallouts indicated that salt stress of more than 8 dSm⁻¹ level strongly inhibited seed germination and plant growth of finger millet and little



millet. However, metamorphosis in salt tolerance among crops may also occur at different growth stages, thus it should be evaluated for different growth stages. Further, at field situation these crops can be tested for response of various levels of saline water irrigation at different growth stages.

Considering the benefit of conservation tillage in rice based cropping scheme a field experiment was carried out to evaluate the impact of conservation tillage on soil health in coastal region of West Bengal during 2011. The design of experiment was split-split plot with cropping scheme (rice-rice and rice-cotton) (kharif—rabi) as main plot treatments and tillage type such as zero tillage (ZT), reduced tillage (RT), and conventional tillage (CT) as sub plot treatments. The residue (R) and no residue (NR) were as sub-sub plot treatments. The soil properties analysed after kharif harvest indicated that salinity was reduced considerably during kharif 5th National Seminar—Climate Resilient Saline Agriculture: Sustaining Livelihood Security 2017

cultivation. EC was slightly higher under rice-cotton scheme than rice-rice scheme. Exchangeable Mg was slightly more than exchangeable Ca. Exchangeable Na was relatively more in cotton-rice scheme than rice-rice scheme. The salinity probe was used to evaluate the bulk soil salinity during the month of May. The rice-rice scheme showed lower bulk soil EC than rice-cotton scheme particularly at 0-30 cm and 30-60 cm soil depth. Also zero tillage plots at surface depth showed higher bulk soil EC than other tillage treatments might be because of higher capillary rise of saline ground water at surface depth in zero tilled plot than other treatments.

There was reduction in bulk density and increase in organic C in ZT than other treatments in surface depth. The soil organic C stock was determined up to 45 cm soil depth and it was highest in RT with residue followed by CT with residue, ZT with residue, RT without residue, CT without residue and lowest in ZT without residue treatment. Organic C stock was more in rice-rice scheme than rice-cotton scheme. The total quantity of soil organic C sequestered within four years of experiment varied from -1.51 to 5.05 Mg C /ha and was linearly related with cumulative C inputs to the soils. The fallouts indicated for sustenance of SOC level (zero change due to cropping) a minimum quantity of 1.86 Mg C /year is required to be added per hectare as inputs. Treatment-wise fraction of soil organic C like very labile, labile, less labile and nonlabile C determined at different concentration of H₂SO₄ and microbial biomass C were also analyzed and passive pool of soil C fraction was more than active pool in treatments when crop residue was added. After four years of experiment in zero tillage, there was 12-18% reduction in yield than other treatments. In Preliminary period of experiment, the yield reduction was up to 28% in case of zero tillage treatment than other treatments.

Agriculture is a backbone of the Indian economy and the rice–wheat cropping scheme is the rib of backbone, but overexploitation of ground water and other available natural resources has fallouted increasing scarcity of resources (water, labour and energy), higher cost of production, diminishing factor productivity, deteriorating soil health shrinking land with emerging concerns of climate change will be a much more challenging task than ever before. These new emerging issues have put a big question mark on the sustainability of rice–wheat cropping scheme. Climate smart agriculture (CSA) offers an attractive alternative to sustainable productivity with efficient use of available resources. Therefore, A participatory strategic research was conducted during, 2014–2016 at farmers' fields in three different climates smart villages viz. Birnarayana, Anjanthali and Chandsamandof Karnal, Haryana,



India under CIMMYT-CCAFS program with the objective of developing and validating portfolios of climate smart agriculture practices in a rice-wheat rotation of Western IGP to provide options and strategies for sustaining productivity and ensure food security in the face of climate change. Six scenarios (S) were established with the various layering of CSAPs including water smart, energy smart, weather smart, carbon smart, nutrient smart and knowledge smart and farmer practices scenario (S1) was used as the base line to compare different sets of CSAP as the five scenarios. Fallout showed that in 2015-16 high magnitude of CSAP helps in maximizing crop productivity (1015%) and profitability (20-25%) while minimizing the adverse effects of associated climatic risks by improving adaptive capacity and reducing mitigation potential of GHG (25-30%) compared to farmer practices scenario (S1). Our study fallouts indicates that the layering of CSA practices have additive effects in terms of improving productivity through better adaptation and also minimizing environmental footprints.

The status of fluoride in groundwater of Ladnu tehsil (Nagaur district) and its effect on wheat and mustard crops were studied for two consecutive years. From the survey area, 100 ground water and soil samples were collected from 31 fluoride affected villages and analysed for fluoride content and categorised in five categories, viz . F1: < 2.0 mg L⁻¹ fluoride, F2: 2.1–4.0 mg L⁻¹ fluoride, F3: 4.1–6.0 mg L⁻¹ fluoride, F4: 6.1–8.0 mg L⁻¹ fluoride and F5: > 8.0 mg L⁻¹ fluoride. Thereafter, for each category of fluoride water four sites were

5th National Seminar–Climate Resilient Saline Agriculture: Sustaining Livelihood Security 2017 i.e . mustard and wheat and also in the quality of groundwater, as a fallout of use of fluoride waters. It was found that 99% of the samples were only marginally alkaline. 23% were in saline category and the rest 67% varied from slightly to moderately saline. Most of the waters had low to moderate sodicity hazard; only 27% waters had high to very high sodicity hazard. Waters of study area also have non alkali to medium alkali hazard and majority of waters falls in low alkali hazard category. Majority of the studied soils were marginally alkaline. Most of the soils had low to moderate sodicity hazard and only 3% represented high sodicity hazard. Fluoride content of groundwater varied from 1.60 to 8.80 mg L⁻¹, indicating a good degree of variation. Further, most of the waters (69%) of the study area had fluoride concentration from 2.1 mg L⁻¹ to 6.0 mg L⁻¹. Positive correlations between soil fluoride and soil pH, soil fluoride and EC were observed indicating that fluoride content increases with soil pH and EC, respectively. While, positive and moderate connection was observed between soil fluoride and sodium. Soil fluoride vs pH of water and soil fluoride vs EC of water also showed positive correlation. The study area lies in semi-arid climate and soils being alkaline have shown less build-up or adsorption of fluoride in soil. Further, the fluoride build-up in soil was in accordance to fluoride concentration in irrigation water and the average build-up in mustard crop was much less than wheat crop. With the increase in fluoride content of water and soil, an increase in fluoride content of plant was also observed, which fallouted in decreased chlorophyll content of leaves, reduction in some plant nutrients (P, K) and thereby reduced crop yield. A highly significant correlation of soil and groundwater fluoride with plant fluoride indicated that both soil and water were the cause for fluoride accumulation in plant. However, accumulation of fluoride in plant varied with plant species i.e. more accumulation in wheat than mustard. Further, as expected much more fluoride content was observed in straw/stover as compared to grains/seed.



Open glowing irrigation in arid parts of Rajasthan: Adaptation to multiple stressors affecting agricultural sustainability Salinity has been a major factor constraining agricultural production in Rajasthan. Selection of salt tolerant varieties (STVs) is considered as one of the environmentally benign and cost effective tactics to overcome the salinity snag. Attempt was made to demonstrate potential of salt tolerant wheat varieties (KRL-210 and KRL- 19) in salt-affected lands of Marwar area of Rajasthan. Study was conducted with 120 farmers from Pali district of Rajasthan. Data were collected using personal interview of the farmers using a glowing structured interview schedule. It was found that all the farmers settled that cultivation of STVs helped increase their income. In terms of social benefits, about 78.5% of the sampled farmers settled that cultivation of STVs helped in the upliftment of small and marginal farmers and in achieving the household food security. In terms of environmental benefits, 96% of the farmers settled that saline land and moderate saline ground water could be used for the cultivation of STVs. Majority of the farmers opined that STVs had more number of tillers (99.6%) and less lodging and shattering tendency (94.7%). All the farmers settled that eating quality of salt tolerant wheat varieties was good. Despite these benefits, there is still a need to create awareness among the farmers in salt-affected areas of Rajasthan about the potential of salt tolerant cultivars in productivity and income enhancements.

Importance and Methodology

1. Intuitive study on Ground-water.
2. Prevention from the disease due to loss of Nutrient in water.
3. Check Mixed Chemicals in Ground Water.
4. Significance study collection of data by hypothetical tests with comparative study through collection of samples from various places.
5. To study the various applications of impurities of water causes diseases and the due to loss of nutrients.
6. Model can help to study the everywhere to prevention of loss of nutrients flow of every kilometer.

We shall use the following methods to complete our Research work:

(i) Charts

- Frequency Polygon
- Bar Graph

(ii) Hypothetical Testing

- Parametric Tests
 - F- Test
 - χ^2 Test
- Non Parametric Tests
 - Rank Test
 - Run Test

(iii) Coefficient of Correlation between

- Himachal Pradesh – Punjab data
- Himachal Pradesh – Rajasthan data
- Punjab – Rajasthan data

(iv) Line of Regression or Curve Fitting

- X on Y
- Y on X
- Exponential Curve
- Power Curve

(v) The Graphs and Charts

- Median Graph (Ogive)
- Regression Graph
- Scatter Diagram

(vi) Mathematical Modelling

- Partial Differential Reckoning
- Solution of Partial Differential Reckoning.
- Standard Deviation Method
- Perturbation Reckoning

Research Work

Nutrients in Water

WHAT ARE NUTRIENTS?


Nutrients, specifically **nitrogen** and **phosphorus**, are chemicals that are essential for plant growth. We add nutrients when we fertilize our gardens and fields, and in the same way, adding nutrients to water fertilizes water-dwelling plants. Nutrients customarily occur at very low concentrations relative to plant demands. Nutrient levels change throughout the year as growing plants take up the nutrients and dying plants release them back into the water.

WHY CARE ABOUT NUTRIENTS



When waterways become over fertilized with nitrogen or phosphorus, heavy plant growth can occur. Excessive plant growth can decrease the aesthetic value of the water because of the smelly decomposing mats of vegetation, and it can create algal blooms which can be toxic. Also, when bacteria decompose dead plant material they use up dissolved oxygen which is important for the survival of macroinvertebrates and other aquatic organisms. If a waterbody doesn't have enough of a nutrient to sustain plant growth, then the nutrient in short supply is called the limiting nutrient.

WHAT IS NITROGEN?



NITROGEN FACTSHEET

Two main groups of nitrogen exist, organic and inorganic. Organic nitrogen includes all of the nitrogen that is part of living animals, animal wastes and the remains of living things. Organic forms of nitrogen must be broken down into inorganic forms in order to be used by plants. Examples of inorganic nitrogen are N_2 , NO_3 , NH_3 , NO_2 . Nitrate (NO_3) is the most onjoint form of inorganic nitrogen found in waterways. Plants can directly use this form of nitrogen to build proteins.

WHY CARE ABOUT NITROGEN?

Environment



When waterways become over fertilized with nitrogen, heavy plant growth can occur. Excessive plant growth can decrease the aesthetic value of the water because of the smelly decomposing mats of vegetation. Also, when bacteria decompose dead plant material they use up dissolved oxygen which is important for the survival of macroinvertebrates and other aquatic organisms.

Human/Animal Health



Nitrates are odorless, colorless, and tasteless so it is important to test feed and drinking water to determine levels of nitrate. High concentrations of nitrate in drinking water can cause methemoglobinemia (also notorious as blue baby syndrome). Concentrations greater than 10 parts per million can be harmful to young babies, and should be avoided by nursing mothers. Concentrations of nitrate over 100 parts per million are toxic to livestock Find out more about nitrate.

Utah Nutrient Standards

- Maximum concentration of Nitrate in drinking water: 10 mg/L
- Greater than 4 mg/L in surface water indicates pollution

NATURAL FACTORS INFLUENCING NITROGEN



- **Runoff-** In Utah concentrations are customarily highest in the springtime when runoff from melting snow carries nutrients from lawns, farms and other areas into the water. Many people in Utah get their culinary water from groundwater from cities or private glowings. Groundwater has naturally higher concentrations of nitrate.
- **Plant uptake-** during the spring and summer months plants grow causing concentrations of nitrates to be low during this time. In the winter and fall, when plants stop growing and die, much of the nitrogen is released into the water again, increasing the nitrogen concentration.

HUMAN FACTORS INFLUENCING NITROGEN





- **Fertilizers-** Because plants require nitrogen, farmers often add nitrogen in the form of fertilizers, sometimes the nitrate leaches into groundwater schemes. Fertilizers can also be washed into surface waters and increase productivity, causing eutrophication.
- **Livestock manure-** Fecal matter contains nitrogen, so when livestock manure washes into surface water (among other pollutant snags), the excess nitrogen can cause eutrophication.
- **Malfunctioning septic schemes-** Similar to livestock manure, when a septic scheme leaches into the ground the bacteria and nutrients can get into ground water schemes and surface water schemes.



- **Discharge from sewage facilities acid precipitation-** Acid precipitation can cause acidification in a water body. Acidification, like eutrophication can lead to a decreased diversity of aquatic species.
- **Resources to limit human influences:** Protect Your Water

Conclusion

Fallouts:

As demand for water increases, water managers and planners need to look widely for ways to improve water management and augment water supplies. The Committee on Ground Water Recharge concludes that artificial recharge can be one option in an integrated strategy to optimize total water resource management, and it believes that with pretreatment, soil-aquifer treatment, and posttreatment as appropriate for the source and site, impaired-quality water can be used as a source for artificial recharge of ground water aquifers.

Artificial recharge using source waters of impaired quality is a sound option where recharge is intended to control saltwater intrusion, reduce land subsidence, maintain stream baseflows, or similar in-ground utilities. It is particularly glowing suited for nonpotable purposes, such as landscape irrigation, because health risks are minimal and public acceptance is high. Where the recharged water is to be used for potable purposes, the health risks and uncertainties are greater. In the past, the development of potable supplies has been guided by the principle that water supply should be taken from the most desirable source feasible, and the rationale for this dictate remains valid. Thus, although indirect potable reuse occurs throughout the nation and world wherever treated wastewater is discharged into a water course or underground and withdrawn downstream or downgradient for potable purposes, such sources are in general less desirable than using a higher quality source for potable purposes. However, when higher-quality, economically feasible sources are unavailable or insufficient, artificially recharged ground water may be an alternative for potable use.

Future Scope:

Ecological and socioeconomic aspects of subterranean hydroschemes have changed during the past 40–50 years. The major environmental pressures (mainly anthropogenic ones) impact the quantity and quality of groundwater resources and the state of subsurface ecosystems, and it is expected that the environmental pressures on groundwater will continue, at least until 2025, unless new environmental policies change this state of affairs. The world demographic increase and the general rise of water demand constitute one of the major environmental pressures on



groundwater ecoschemes especially in less developed countries in Africa, Asia and South America. Specific human activities leading to the depletion of groundwater reserves include agricultural practices, landscape alteration, urbanization demand for domestic and public drinking water, various industrial activities such as thermoelectric production and mining, and the rise of tourism in coastal areas. Climate change is contributing to the water crisis too, especially in areas with arid climate and/or in some humid monsoonal countries. The overload of aquifers with pollutants derived from agriculture (fertilizers and pesticides), from industry (release of hydrocarbon chemicals, especially spills), from waste and industrial waters, from domestic and industrial landfills, from the infiltration of pollutants from surface and from the intrusion of saline water affect groundwater quality. The dangerous increase in contaminated subsurface sites with chemicals and microbial pathogens brings with it health risks to humans. Changes of redox situation in groundwater zones, changes of biological diversity, vegetation changes with modification of agriculture practices and impacts at the biosphere scale, such as the increase in the concentration of nitrous oxides in the atmosphere, all impact groundwater ecoschemes. Groundwater ecoschemes must be better investigated and understood. Economic, social and ecological lines of thinking have to be combined in order to achieve meaningful policies for the sustainable development of groundwater reserves and for the protection of subsurface ecoschemes. Practical measures and ideas for the development of policies up to the 2025 time-horizon should improve the sustainable usage of the world's groundwater resources.

Bibliography

1. Andreski, Stanislav (1972). Social Sciences as Sorcery. St. Martin's Press. ISBN 0-14-021816-5.
2. Truesdell, Clifford (1984). An Idiot's Fugitive Essays on Science. Springer. pp. 121–7. ISBN 3-540-90703-3.
3. Billings S.A. (2013), Non-linear Scheme Identification: NARMAX Methods in the Time, Frequency, and Spatio-Temporal Domains, Wiley.
4. "Thomas Kuhn". Stanford Encyclopedia of Philosophy. 13 August 2004. Retrieved 15 January 2019.
5. Thornton, Chris. "Machine Learning Lecture". Retrieved 2019-02-06.
6. Pyke, G. H. (1984). "Optimal Foraging Theory: A Critical Review". Annual Review of Ecology and Systematics. 15: 523–575. doi:10.1146/annurev.es.15.110184.002515.
7. "GIS Delineations of Terminology M-P". LAND INFO Worldwide Mapping. Retrieved January 27, 2020.
8. Gallistel (1990). The Organization of Learning. Cambridge: The MIT Press. ISBN 0-262-07113-4.
9. Whishaw, I. Q.; Hines, D. J.; Wallace, D. G. (2001). "Dead reckoning (path integration) requires the hippocampal formation: Evidence from spontaneous exploration and spatial learning tasks in light (allothetic) and dark (idiothetic) tests".
10. Behavioural Brain Research. 127 (1–2): 49–69. doi:10.1016/S0166-4328(01)00359-X. PMID 11718884.
11. "Archived copy". Archived from the original on 19 January 2012. Retrieved 19 January 2012.
12. "Temperature Regimes". Water Resource Department - Government Of Rajasthan. Retrieved 6 December 2008.
- INDIA METEOROLOGICAL DEPT
13. "District Census 2011". Census2011.co.in. 2011. Retrieved 30 September 2011.