

Multivalent Uniformly Convex Functions by Using Differential Operator

Pooja Lakhera, Research Scholar, Department of Mathematics, SunRise University, Alwar.

Dr. Rajeev Kumar, Associate Professor, Department of Mathematics, SunRise University, Alwar.

This section is devoted to regular and multi-valent mapping by using differential Op_{tor} in the \mathcal{U}_D . We study various exciting things for this novel class prior to multivalent mappings.

Allow S exist the class of the mappings

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n, \quad a^n \geq 0 \quad (p \in N) \quad (1.1)$$

whichever regular & p - valent within effective \mathcal{U}_D , $U = \{z: |z| < 1\}$

Furthermore S^* be effective sub \mathcal{CL}_{ss} prior to S containing to mappings

$$f(z) = z^p - \sum_{n=p+1}^{\infty} a_n z^n, \quad a^n \geq 0 \quad (p \in N) \quad (1.2)$$

$$\operatorname{Real} \left\{ 1 + \frac{zf''(z)}{f'(z)} - \alpha \right\} > \beta \left| \frac{zf'(z)}{f(z)} - 1 \right|, \quad (z \in U) \quad (1.3)$$

Wherever $-1\alpha \leq 1$, $\beta \geq 0$ & $p \in N$

(ii) A mappings $f(z) \in S$ suppose to subsist within effective \mathcal{CL}_{ss} $\mathcal{UCV}(\alpha, \beta)$ to consistently β - \mathcal{CV} & satisfy

$$\operatorname{Real} \left\{ 1 + \frac{zf''(z)}{f'(z)} - \alpha \right\} > \beta \left| \frac{zf''(z)}{f'(z)} - 1 \right|, \quad (z \in U) \quad (1.4)$$

wherever $\alpha \leq 1$, $\beta > 0$ and $p \in N$

From above (1.3) & (1.4)

$f(z) \in \mathcal{UCN}(\alpha, \beta)$ do comparable toward $zf'(z) \in S_p(\alpha, \beta)$

Hd_{pro} of $f(z)$, $g(z) \in S$ can be define as

$$f * g(z) = z^p + \sum_{k=p+1}^{\infty} a_k b_k z^k \quad (z \in U), \quad p \in N \quad (1.6)$$

Concerning effective mapping $f(z) \in S$, without help classify affecting subsequent

$$I^0 f(z) = f(z), \quad I^1 f(z) = zf'(z) + \frac{1+p}{z^p}$$

along with $k = 2, 3, 4, \dots$

$$= z^p + \sum_{n=p+1}^{\infty} n(k) a_n z^n, \quad p \in N \quad (1.7)$$

Somewhere I^k is the same as diff. Op_{tor} , Ghanim & Darus [2], S.K.Lee,

S. Khairnar with S. Rajas [9] have studied this Op_{tor} widely.

Let $S^*(\alpha, \beta) \in S$ consisting of the mapping of the form (1.1) and satisfy

$$\left| \frac{\frac{z(I^k f(z))' - p}{I^k f(z)} - \mu}{\frac{\beta z(I^k f(z))' - \alpha p z}{I^k f(z)} - \alpha p z} \right| < \mu \quad (1.8)$$

where $-1 \leq \alpha < \beta \leq 1$ and $0 < \mu \leq 1$ ($z \in U$).

Also let $S^{**}(\alpha, \beta) = S^*(\alpha, \beta) \cap S^*$

3.2.1 Coefficient Estimate

Here we obtained a essential & enough situation for function $f(z)$ inside effective \mathcal{CL}_{ss} $S^*(\alpha, \beta)$ and $S^{**}(\alpha, \beta)$.

Theorem 1: A mapping of the equation (1.1) is in $S^*(\alpha, \beta)$ iff

$$\sum_{n=1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| < \mu p(\beta - \alpha p), \quad (1.9)$$

where $-1 \leq \alpha < \beta \leq 1$ and $0 < \mu \leq 1$ and $p \in N$.

Proof : It's enough to illustrate so as to

$$\left| \frac{\frac{z(I^k f(z))' - p}{I^k f(z)}}{\frac{\beta z(I^k f(z))' - \alpha p z}{I^k f(z)}} \right| < \mu$$

as $f(z) \in S^*(\alpha, \beta)$ we have

$$\begin{aligned} & \left| \frac{\frac{z(I^k f(z))' - p}{I^k f(z)}}{\frac{\beta z(I^k f(z))' - \alpha p z}{I^k f(z)}} \right| \leq \mu (z \in \mathcal{U}, p \in N) \\ & = \left| \frac{\frac{pz^p + \sum_{n=p+1}^{\infty} n(k)na_n z^n}{z^p + \sum_{n=p+1}^{\infty} n(k)a_n z^n} - p}{\frac{\beta pz^p + \beta \sum_{n=p+1}^{\infty} n(k)na_n z^n}{z^p + \sum_{n=p+1}^{\infty} n(k)a_n z^n} - \alpha p} \right| \leq \mu \\ & = \left| \frac{pz^p + \sum_{n=p+1}^{\infty} n(k)na_n z^n - pz^p - \sum_{n=p+1}^{\infty} n(k)a_n z^n}{pz^p(\beta - \alpha p) + \sum_{n=p+1}^{\infty} (\beta - \alpha p)n(k)na_n z^n} \right| \leq \mu \end{aligned}$$

$$\sum_{n=1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| |z^n| < \mu p(\beta - \alpha p) |z^p|$$

Allowing the value of $z \rightarrow -1$ taking place effective $\mathcal{R}e_{al} A_{is}$, without help obtained

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| < \mu p(\beta - \alpha p)$$

Theorem 2: A essential and enough stipulation in favor of $f(z)$ prior to the structure (1.2) toward exist effective $\mathcal{C}\ell_{ss}$ $S^{**}(\alpha, \beta)$.

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| \leq \mu p(\beta - \alpha p) \quad (1.10)$$

where $-1 \leq \alpha \leq \beta$ and $0 < \mu \leq 1$ & $p \in N$.

Proof : It's enough to illustrate so as to

$$\left| \frac{\frac{z(I^k f(z))' - p}{I^k f(z)}}{\frac{\beta z(I^k f(z))'}{I^k f(z)} - \alpha p \frac{z(I^k f(z))}{I^k f(z)}} \right| \leq \mu$$

We enclose

$$\left| \frac{\frac{pz^p + \sum_{n=p+1}^{\infty} n(k)na_n z^n}{z^p + \sum_{n=p+1}^{\infty} n(k)a_n z^n} - p}{\frac{\beta pz^p + \beta \sum_{n=p+1}^{\infty} n(k)na_n z^n}{z^p + \sum_{n=p+1}^{\infty} n(k)a_n z^n} - \alpha p} \right| \leq \mu$$

$$\sum_{n=1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| |z^n| \leq \mu p(\beta - \alpha p) |z^n|$$

Allowing the value of $z \rightarrow -1$ with effective $\mathcal{R}e_{al} Ax_{is}$, without help acquire

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| < \mu p(\beta - \alpha p)$$

The $S^{**}(\alpha, \beta)$ remain closed underneath linear combination we will prove this in the following theorem.

Theorem 3: If $f(z)$ is definite through (1.2) and

$$g(z) = z^p - \sum_{n=p+1}^{\infty} b_n z^n$$

live in the class $S^{**}(\alpha, \beta)$. Then the function

$$h(z) = (1 - \dot{\alpha}) f(z) + \dot{\alpha} g(z) = z^p - \sum_{n=p+1}^{\infty} \eta_n z^n \quad (1.11)$$

Is as well within $S^{**}(\alpha, \beta)$ wherever

$$\eta_n = (1 - \epsilon) a_n + \epsilon b_n \quad 0 \leq \epsilon \leq 1.$$

Proof : As the mappings $f(z)$ & $g(z)$ hold inside $S^{**}(\alpha, \beta)$, so we include

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| < \mu p(\beta - \alpha p)$$

And

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |b_n| < \mu p(\beta - \alpha p)$$

Then

$$\begin{aligned} h(z) &= (1 - \epsilon) f(z) + \epsilon g(z) \\ &= (1 - \dot{\alpha}) z - \sum_{n=p+1}^{\infty} a_n z^n + \dot{\alpha} \left(z - \sum_{n=p+1}^{\infty} b_n z^n \right) \\ &= z^p - \sum_{n=p+1}^{\infty} \left[(1 - \dot{\alpha}) a_n + \dot{\alpha} b_n \right] z^n \end{aligned}$$

$$= z^p - \sum_{n=p+1}^{\infty} c_n z^n$$

when $c_n = (1 - \epsilon)a_n + \epsilon b_n$

Now consider

$$\begin{aligned} & \sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |c_n| \\ &= \sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |(1 - \epsilon)a_n + \epsilon b_n| \\ &\leq (1 - \delta) \sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| \\ &\quad + \delta \sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| \\ &\leq (1 - \epsilon)\mu p(\beta - \alpha p) + \epsilon\mu p(\beta - \alpha p) \\ &= \mu p(\beta - \alpha p) \end{aligned}$$

Thus we get

$$\sum_{n=p+1}^{\infty} [(n-p) + \mu n(\beta - \alpha p)] n(k) |a_n| < \mu p(\beta - \alpha p)$$

Hence $h(z) \in S^{**}(\alpha, \beta)$

References :

1. Aghalary R. and Kulkarni S.R. (2002), Some theorems on univalent functions, *J. Indian Acad. Math.*, 24(1), 81-93.
2. Ali Muhammad (2012), On some applications of subordination and superordination of multivalent functions involving the extended fractional differintegral operator, *Le Matematiche* Vol. LXVII– Fasc. II, pp. 59–75.
3. Amer A. and Darus M., (2012), On a subclass of k uniformly starlike functions associated with the generalized hypergeometric functions, *Adv. Studies Theor. Phys.*, Vol.6, 273-284.
4. Aqlan E., Jahangiri J.M. and Kulkarni S.R. (2004), Classes of K -uniformly convex and starlike functions, *Tamkang J. Math.*, 35(3), 261-266.
5. Asthan W.G., Mustafa H.D and Mouajeeb E.K. (2013), Subclass Of multivalent functions defined by Hadamard product involving a linear operator, *Int. Journal of Math. Analysis*, 7, 24, 1193-1206.
6. Athsan W. G. and Kulkarni S.R., (2008), Generalized Ruscheweyh derivatives involving a general fractional derivative operator defined on a class of multivalent functions II, *Int. Journal Of Math. Analysis*, Vol.2,no. 3, 97-109.
7. Athsan, W.G. And Kulkarni S.R.,(2007), On a Class of p -Valent Meromorphic Functions Defined by Integral Operator, *International J. Of Math. Sci. & Engg. Appl.* 1, 129-140.
8. Athsan, W.G. and Kulkarni, S.R.(2008), New Classes of Multivalently Harmonic Functions, *Int. journal of Math. Analysis*, 2, 3, 111-121.
9. B.A. Frasin and M. Darus (2004), Integral means and neighborhoods for analytic univalent functions with negative coefficients, *Soochow Journal of Mathematics* , Vol.30 No. 2 , 217-223.

